



Excited state properties of a short π -electron conjugated peridinin analogue



Nikki M. Magdaong^a, Dariusz M. Niedzwiedzki^b, Jordan A. Greco^a, Hongbin Liu^a, Koki Yano^c, Takayuki Kajikawa^c, Kazuhiko Sakaguchi^d, Shigeo Katsumura^{a,e}, Robert R. Birge^{a,*}, Harry A. Frank^{a,*}

^a Department of Chemistry, University of Connecticut, Storrs, CT 06269-3060, USA

^b Photosynthetic Antenna Research Center, Washington University in St. Louis, MO 63130, USA

^c Department of Chemistry, School of Science and Technology, Kwansei Gakuin University, Gakuen, Sanda, Hyogo 669-1337, Japan

^d Graduate School of Science, Osaka City University, 3-3-138 Sugimoto, Sumiyoshi-ku, Osaka 558-8585, Japan

^e Osaka City University Advanced Research Institute for Natural Science and Technology, 3-3-138 Sugimoto, Sumiyoshi-ku, Osaka 558-8585, Japan

ARTICLE INFO

Article history:

Received 12 November 2013

In final form 3 January 2014

Available online 8 January 2014

ABSTRACT

C_{29} -peridinin is a synthetic analogue of the important, naturally-occurring carotenoid, peridinin, found in several marine algal species. C_{29} -peridinin has five conjugated carbon–carbon double bonds compared to eight possessed by peridinin and also lacks the methyl group functionalities typically present along the polyene chain of carotenoids. These structural modifications lead to unique excited state properties and important insights regarding the factors controlling the photophysics of peridinin and other carbonyl-containing carotenoids, which are critical components of the light-harvesting systems of many photosynthetic organisms.

© 2014 Elsevier B.V. All rights reserved.

1. Introduction

Ocean-dwelling algal organisms are well-known for the proliferation of carbonyl-containing carotenoids in light-harvesting pigment–protein complexes that facilitate capturing sunlight for photosynthetic growth [1–3]. These functionally derivatized keto-carotenoids exhibit spectroscopic and kinetic behavior that can be strikingly different from carotenoids that lack a carbonyl group in conjugation with the π -electron system of conjugated carbon–carbon double bonds. One of the characteristic differences is that the lifetime of the lowest-lying excited state, S_1 , of carotenoids possessing a carbonyl group in conjugation with the polyene backbone is strongly dependent on the polarity of the solvent [4–8]. This is unusual because the S_1 state of carotenoids is a state into which absorption from the ground state, S_0 , is quantum mechanically forbidden, and therefore the spectra and dynamics associated with S_1 are typically not affected by the solvent environment [9–13]. The forbiddenness of the $S_0 \rightarrow S_1$ transition is due to the fact that both S_0 and S_1 have A_g^- symmetry in the idealized C_{2h} point group. According to the selection rules for one-photon optical transitions, a change in symmetry and pseudoparity is required for the transition to be allowed. The strongly allowed transition that gives carotenoids their vibrant visible coloration occurs between the S_0

($1^1A_g^-$) and S_2 ($1^1B_u^+$) states, which differ in both symmetry and pseudoparity [14].

The dependence of the S_1 lifetime of carbonyl-containing carotenoids has been explained by the formation of an intramolecular charge transfer (ICT) state whose energy and electronic coupling is modulated by the solvent polarity [5]. Recent ultrafast time-resolved spectroscopic and computational investigations of the carbonyl-containing carotenoid, peridinin, have suggested that after photoexcitation into the S_2 ($1^1B_u^+$) state, a shift of electron density from the allenic side of peridinin toward the lactone ring occurs resulting in a bond-order reversal along the polyene chain [15]. These effects are accompanied by solvent reorganization, which together generate the ICT state through quantum mechanical mixing of the S_2 ($1^1B_u^+$) ionic state with the lowest-lying S_1 ($2^1A_g^-$) covalent state. The charge transfer character evolves in less than 100 fs and results in a very large (~ 35 D) dipole moment.

In order to explore the nature of the ICT state, several analogues of peridinin having various extents of π -electron conjugation (Figure 1) have been synthesized and characterized spectroscopically and computationally [16–20]. Naturally-occurring peridinin has a C_{37} carbon skeleton rather than the typical C_{40} system present in most carotenoids [21]. In this Letter new results are presented on the shortest member of this series of synthetic peridinin, an analogue that has a C_{29} carbon skeleton, hereafter denoted C_{29} -peridinin. This molecule is not only a shortened version of peridinin: C_{29} -peridinin has five conjugated carbon–carbon double bonds compared to eight possessed by peridinin (Figure 1), but it also lacks the methyl functionalities typically

* Corresponding authors. Address: Department of Chemistry, 55 North Eagleville Road, University of Connecticut, Storrs, CT 06269-3060, USA. Fax: +1 860 486 6558.

E-mail addresses: rbirge@uconn.edu (R.R. Birge), harry.frank@uconn.edu (H.A. Frank).




Figure 1. Structure of C₂₉-peridinin and other peridinins.

present along the polyene chain of carotenoids. These structural modifications lead to unique excited state spectral and kinetic properties and provide important insights regarding the factors that control the photophysics of peridinin and other carbonyl-containing carotenoids that are critical components in the light-harvesting systems of an abundance of photosynthetic organisms.

2. Materials and methods

2.1. Sample preparation

The details of the synthesis of C₂₉-peridinin will be reported elsewhere. Prior to the optical experiments, the molecule was dissolved in acetonitrile and injected into a Millipore Waters 600E high-performance liquid chromatography (HPLC) employing a C₃₀ YMC column and an isocratic mobile phase protocol consisting of 87:10:3, acetonitrile:methanol:water (v/v/v) at a flow rate of either 0.8 or 1 mL/min. The sample volume was 200 μ L for each injection. Pure C₂₉-peridinin eluting from the column was identified using a Waters 996 single diode-array detector, collected, dried using a gentle stream of gaseous nitrogen, and stored at -80 °C until ready for use.

2.2. Spectroscopic methods

Steady-state absorption and fluorescence: All of the spectroscopic experiments were carried out at room temperature. C₂₉-peridinin was dissolved in spectroscopic grade *n*-hexane

(Sigma Aldrich), tetrahydrofuran (Alfa Aesar), acetonitrile (Sigma Aldrich), and methanol (Sigma Aldrich). Absorption spectra were recorded using a Cary 50 UV-visible spectrometer. Fluorescence spectroscopy was carried out using a Jobin-Yvon Horiba Fluorolog-3 model FL3-22 equipped with double monochromators having 1200 grooves/mm gratings, a Hamamatsu R928P photomultiplier tube detector, and a 450 W ozone-free Osram XBO xenon arc lamp. The emission spectra were collected using 376 nm (*n*-hexane) and 400 nm (tetrahydrofuran, methanol, acetonitrile) excitation at a right angle relative to the emission beam and corrected using a file generated using a 200 W quartz tungsten-halogen filament lamp. Both the emission and excitation slit widths were set to a bandpass value of 3 nm (*n*-hexane, tetrahydrofuran, methanol) or 5 nm (acetonitrile).

Time-resolved fluorescence: Time-resolved fluorescence data were obtained at room temperature using the setup previously described [20]. The samples had an OD of ~ 0.3 at the maximum in their absorption recorded in a 1 cm cuvette. Depending on the solvent, the excitation beam had a wavelength between 380 and 430 nm and was positioned at a right angle relative to the detector. The beam power was 40–130 mW (at 80 MHz frequency) which corresponds to 0.5–1.5 nJ of energy and, with a spot size ~ 0.5 mm in diameter, yielded a photon intensity of $5\text{--}10 \times 10^{11}$ photons/cm².

Transient absorption spectroscopy: Time-resolved pump-probe absorption experiments were performed at room temperature using Helios, a femtosecond transient absorption spectrometer (Ultrafast Systems LLC, Sarasota, FL, USA) coupled to a femtosecond laser system described previously [22]. The samples were excited with pump beam energy of 500 nJ in a spot size of 1 mm diameter corresponding to an intensity of $\sim 2 \times 10^{14}$ photons/cm². The pump wavelength was 400 nm. Steady-state absorption spectra were recorded before and after all transient experiments to confirm the stability of the C₂₉-peridinin sample during laser excitation.

2.3. Computational methods

The theoretical methods used in the present Letter are identical to those used previously by Wagner et al. [15] to investigate the excited state properties of peridinin. The interested reader is directed to that work for a discussion of the methods, procedures, and rationale.

3. Results and discussion

To better understand the nature of the steady-state absorption and fluorescence spectral features of C₂₉-peridinin, it is useful to view them in the same context as the spectra recorded from peridinin and the C₃₃-, C₃₅- and C₃₉-peridinin analogues in the same solvents. The steady-state absorption and fluorescence spectra recorded at room temperature in *n*-hexane and methanol from all of these molecules are shown in Figure 2.

The absorption spectrum of C₂₉-peridinin, which corresponds to an S₀ \rightarrow S₂ transition, is significantly blue-shifted relative to the spectra from the other molecules. In *n*-hexane, the spectral origin (0–0) vibronic band of C₂₉-peridinin is sufficiently resolved that it can be located at ~ 400 nm; i.e., the S₀ \rightarrow S₂ transition corresponds to an energy of ~ 25000 cm⁻¹. As the solvent polarity increases, the absorption band broadens substantially, but retains a small amount of vibronic structure compared to the other peridinins, as can be seen in the absorption spectrum of C₂₉-peridinin in methanol (bottom left panel of Figure 2).

Room temperature fluorescence spectra of C₂₉-peridinin in *n*-hexane and methanol are also shown in Figure 2 along with the spectra from peridinin and the other peridinin analogues. The




Figure 2. Steady-state absorption and emission spectra. Data for peridinin, C_{33} -, C_{35} -, and C_{39} -peridinin in *n*-hexane and methanol were taken from Ref. [17].

spectra are broad and red-shifted compared to their respective absorption spectra in both solvents, but clearly resemble fluorescence emission lineshapes associated with the $S_1 \rightarrow S_0$ transition. An exception is C_{39} -peridinin, which displays emission from both the S_1 state, whose (0–0) band is at ~ 650 nm, and the S_2 state, which is characterized by an emission band at ~ 540 nm in both solvents.

As the π -electron conjugated chain length decreases, the energy difference between the S_2 and S_1 states also decreases. For the longest molecule in the series, C_{39} -peridinin, the energy difference between the S_2 and S_1 states is ~ 3100 cm^{-1} based on the position of their (0–0) absorption and fluorescence bands. For C_{29} -peridinin, the (0–0) band of the S_1 fluorescence occurs at ~ 410 nm which is very close to the (0–0) band of the absorption spectrum at ~ 400 nm, resulting in a substantially smaller S_2 – S_1 energy difference of ~ 600 cm^{-1} . Also, the intensity of emission decreases substantially as the solvent polarity increases. The relative fluorescence quantum yield of C_{29} -peridinin is smaller by approximately an order of magnitude in methanol compared to in the nonpolar solvent, *n*-hexane.

The transient absorption spectra of C_{29} -peridinin recorded in *n*-hexane, tetrahydrofuran, methanol, and acetonitrile in the visible spectral region at various delay times after excitation into the S_2 state are shown in the first column of Figure 3A. As indicated above for the steady-state spectra, in order to understand the nature of the transient absorption spectral bands, it is useful to view them in context with the transient absorption spectra from peridinin and the other peridinin analogues recorded in the same solvents. These spectra are shown in Figure 3A. Note that excitation of the longest molecule in the series, C_{39} -peridinin dissolved in the non-polar solvent, *n*-hexane (top panel in the last column of Figure 3A), results in a rapid bleaching of the $S_0 \rightarrow S_2$ absorption bands followed in a few hundred femtoseconds by the build-up of a narrow excited state absorption (ESA) band at ~ 540 nm. As the polarity of the solvent is increased, an additional broader ESA band appears between 550 and 700 nm concomitant with the narrow band. However, unlike the narrow ESA band whose position remains relatively unaffected by solvent polarity, the broader feature in the spectra from C_{39} -peridinin increases in intensity and shifts from longer to shorter wavelength with increasing solvent polarity. This suggests that the narrow and broad bands belong to two different electronic transitions.

As the π -electron conjugation is shortened by going down the series from C_{39} -peridinin to peridinin, C_{35} -peridinin, C_{33} -peridinin and C_{29} -peridinin, both the broad and narrow ESA spectral bands shift to shorter wavelength owing to a decrease in the extent of π -electron conjugation (top row of spectra in Figure 3A). In *n*-hexane, the narrow ESA band shifts ~ 40 nm to the blue for each one fewer double bond. Also, in this solvent, as the π -electron conjugation length of the molecule is decreased, the intensity of the narrow ESA band becomes less pronounced relative to that of the broad band, which for C_{29} -peridinin in *n*-hexane, appears at ~ 560 nm (top left hand traces in Figure 3A). The narrow ESA band in the transient absorption spectra of C_{29} -peridinin recorded in *n*-hexane has shifted to such a short wavelength (< 440 nm) that it is out of the spectral response window of the spectrometer. Also for C_{29} -peridinin, another broad band not observed in the transient absorption spectra of any of the other peridinin appears at ~ 700 nm, and its wavelength remains relatively invariant to solvent polarity as seen in the ESA spectra shown in the first column of Figure 3A. Like the broad band at ~ 560 nm, this additional band is prominently observed in the ESA traces taken at the shortest (< 150 fs) delay times. This indicates that the state from which these two broad ESA transitions originate is populated from the S_2 state. At subsequent times, both broad bands (the one at ~ 560 nm in *n*-hexane that is shifted to the blue with solvent polarity and the one at ~ 700 nm that does not shift) decay at the same rate. This suggests that the states from which these two spectral features originate are in fast dynamic equilibrium.

The transient absorption spectra of the peridinin in the different solvents recorded in the near infrared (NIR) spectral region at various delay times after excitation into its S_2 state are shown in Figure 3B. Focusing initially on C_{39} -peridinin (last column in Figure 3B), there is a strong band at ~ 900 nm that rises and decays within the time course of the laser excitation pulse indicating it is an ESA transition originating from the S_2 state. This band is not affected by solvent polarity, but shifts to shorter wavelength upon decreasing the π -electron conjugation length of the molecule. This is seen most clearly in the ESA spectra for the series of peridinin recorded in *n*-hexane (top row of spectra in Figure 3B). As the fast feature decays, a lower-energy band appears at ~ 1450 nm for C_{39} -peridinin in *n*-hexane, which suggests that it is directly populated from the S_2 state subsequent to photoexcitation. This band decays in several picoseconds depending on the solvent,




Figure 3. Transient absorption spectra recorded in the (A) visible and (B) NIR spectral regions. Data for peridinin, C₃₃-, C₃₅-, and C₃₉-peridinin in *n*-hexane, tetrahydrofuran (THF), and acetonitrile (ACN) were taken from reference [20], while data for C₃₅-, peridinin and C₃₉-peridinin in methanol were taken from Ref. [17].

and shifts slightly to shorter wavelength and becomes stronger as the π -electron conjugation of the molecule is decreased; e.g., see top row of spectra in Figure 3B. The band appears at ~ 1350 nm for C_{29} -peridinin. This longer-lived band also becomes broader and shifts to the blue with increasing solvent polarity; e.g., see the first column of spectra in Figure 3B. The fact that this band is long-lived suggests that it originates from the lowest excited state of the molecules. For peridinin and C_{39} -peridinin, this band is very likely a higher vibronic feature associated with the $S_1 \rightarrow S_2$ transition as suggested by Zigmantas et al. [7]. However, this feature cannot be an $S_1 \rightarrow S_2$ transition for the shortest in the series, C_{29} -peridinin, because the (0–0) vibronic bands of the $S_0 \rightarrow S_2$ absorption and $S_2 \rightarrow S_0$ fluorescence spectra differ by only ~ 10 nm. This indicates an energy separation between the S_1 and S_2 states of only ~ 600 cm^{-1} (see above), which would put the $S_1 \rightarrow S_2$ transition at a very long wavelength. Therefore, the band must be associated with a transition to a higher-lying state than S_2 .

In addition to the ESA features in the transient spectra, stimulated emission is evident as broad negative amplitude, particularly in the spectra of the molecules dissolved in polar solvents (see for example the transient NIR spectra from C_{35} -peridinin in tetrahydrofuran, methanol, and acetonitrile, center column in Figure 3B). This emission has been assigned as originating from the ICT state [6,7,20,23–25].

In order to probe more deeply into the origin and dynamics of the electronic states giving rise to the transient spectra, global fitting according to a sequential decay model was done on the spectral and temporal datasets from C_{29} -peridinin. The resulting amplitude traces are termed evolution associated decay spectra (EADS) [26] and are shown in Figure 4 for the datasets taken in the visible region. In all solvents, four kinetic components were required to achieve a satisfactory fit based on a chi square (χ^2) test and minimization of the residuals [26,27].




Figure 4. Evolution associated difference spectra (EADS, left column) obtained from the global fitting of the transient absorption in the visible region and time-resolved fluorescence kinetic profiles (TRF, right column) of C_{29} -peridinin in different solvents.

The EADS traces for C_{29} -peridinin in all of the solvents in the visible region (Figure 4) show a very fast 150–170 fs decay component that is very likely associated with the lifetime of the photoexcited S_2 state. The bleaching of the $S_0 \rightarrow S_2$ absorption profile is not evident because that band occurs below 425 nm which is outside the spectral response window of the spectrometer. The initial fast component decays into a second EADS component that has the two broad bands at ~ 560 and ~ 700 nm described above. This EADS component has a lifetime of 6.3 ps in *n*-hexane, 1.3 ps in tetrahydrofuran, 440 fs in methanol and 370 fs in acetonitrile. Except for the molecule dissolved in *n*-hexane, and consistent with previous analyses of the other peridinin analogues [20], this decay component is associated with a vibronically hot S_1 state. This is evidenced by the fact that, for the molecule in the polar solvents, the spectral lineshape narrows and shifts to shorter wavelength as it decays into a third EADS component [27,28]. In *n*-hexane there are no spectral features that can be attributed to vibrational relaxation which suggests that in this solvent, the process is complete prior to deactivation from the S_2 state. The lifetime of the third EADS component decreases with increasing solvent polarity (Table 1) suggestive of different amounts of ICT character in the excited state associated with this component [20]. However, a fourth EADS component is also needed to fit the data which is not the case for any of the longer peridinin analogues [17]. The lifetime of this fourth component also decreases with solvent polarity (Table 1) and is similarly suggestive of different amounts of ICT character in the state. For C_{29} -peridinin in *n*-hexane the fourth EADS component appears very small because the $S_1 \rightarrow S_N$ transition is out of the spectral window of the spectrometer. Because of its small amplitude and apparently lengthy lifetime, the global fitting assigns this component as infinitely long. In actuality, the lifetime of the S_1 state of C_{29} -peridinin in *n*-hexane is 2.9 ns as measured more precisely using time-resolved fluorescence (right hand column in Figure 4). As the solvent polarity is increased, some EADS amplitude can be observed at very short wavelengths, and the lifetime of this fourth EADS becomes shorter at 53 ps in tetrahydrofuran, 17 ps in methanol, and 13.5 ps in acetonitrile. These values are wholly consistent with those measured by time-resolved fluorescence which reveal 81 ps in tetrahydrofuran, 16.5 ps in methanol, and 15 ps in acetonitrile (Figure 4). The fact that a fourth EADS component is necessary at all is highly suggestive that the third and fourth components are associated with different conformations of the molecule having different excited state decay times. This interpretation is consistent with the quantum computational results described below. Table 1 summarizes the kinetics data from the transient absorption and time-resolved fluorescence experiments.

The global fits to the datasets taken in the NIR region are given in the supplemental information (Figure S1) and depending on the solvent, required three or four components for a good fit. The kinetic parameters derived from both the visible and NIR datasets are summarized in Table 1 and show good agreement among the values. However, for the experiments in the more polar solvents, the component assigned to the vibronically hot state from the EADS analysis of the visible datasets is not resolved, presumably because of low amplitude of this component in the NIR spectral region.

From the trends in the positions and amplitudes of the transient spectral features seen for the entire series of peridinin shown in Figure 3A and B, it is clear there are four different spectral bands of C_{29} -peridinin that need explanation: Two of these bands, one in the NIR and one in the visible region, broaden and shift to higher energy with increasing solvent polarity, and two of the bands (at ~ 700 nm and <440 nm) remain relatively unaffected by changing the solvent. These observations suggest that the bands that are affected by the solvent originate from the ICT state, which has significant B_u^+ character [15,17,20,28]. Because this state has a large

Table 1
Dynamics of the excited states of C₂₉-peridinin obtained in various solvents using transient absorption (TA) and time-resolved fluorescence (TRF) spectroscopy.^a

Solvent	Probe region	TA				TRF	
		τ_1	τ_2	τ_3	τ_4	τ_{F1}	τ_{F2}
n-Hexane	Vis	0.15 ± 0.01	6.3 ± 0.2	21 ± 1	inf	<FWHM	2900 ± 100
	NIR	0.24 ± 0.01	6.3 ± 0.1	25 ± 2	inf	(300)	
THF	Vis	0.15 ± 0.01	1.3 ± 0.1	8.2 ± 0.2	53 ± 3	n.e.	81 ± 2
	NIR	0.28 ± 0.2	5.6 ± 0.5	58 ± 3		(28)	
MeOH	Vis	0.17 ± 0.01	0.44 ± 0.02	3.9 ± 0.2	17 ± 1	<FWHM	16.5 ± 0.5
	NIR	0.17 ± 0.01	2.9 ± 0.3	16 ± 1		(12)	
ACN	Vis	0.16 ± 0.01	0.37 ± 0.01	1.9 ± 0.1	13.5 ± 0.4	<FWHM	15.0 ± 0.5
	NIR	0.18 ± 0.01	1.3 ± 0.1	13.6 ± 0.6		(12)	

^a The lifetimes are all given in picosecond units and were obtained from the global fitting of the transient absorption (TA) and time-resolved fluorescence (TRF) datasets. The uncertainties were determined by exploring the region of solution for each fitted parameter based on the goodness of fit and values of the residuals. FWHM – full width at half maximum of the instrument response function (in parenthesis in picoseconds); n.e., not evident, THF, tetrahydrofuran; MeOH, methanol; ACN, acetonitrile.




Figure 5. Effect of solvent environment on the low-lying Franck–Condon (FC) and relaxed (S₁ rlx) excited singlet states of C₂₉-peridinin and peridinin based on EOM-CCSD and PCM procedures. Transition energies are relative to the third-order Møller-Plesset (MP3) ground state. Each excited state is represented by a rectangle, the height of which is proportional to the oscillator strength (marked above or below the rectangle) and the ionic versus covalent character is indicated by color (see insert). All symmetry assignments are approximate based on the symmetry labels of parent linear polyenes.

dipole moment, it is expected to decrease in energy with increasing solvent polarity, thereby leading to a shift to higher energy of the transitions originating from it. The bands that are unaffected by the solvent most likely originate from the S₁ state, which has a significant A_g⁻ symmetry known to be less prone to changing its position upon alteration of the solvent polarity.

As the above discussion suggests, C₂₉-peridinin is somewhat of an outlier in terms of its spectroscopic and photophysical properties compared to the other peridinin derivatives. The two lowest excited states of C₂₉-peridinin are observed to be much closer in energy than for the other peridinin derivatives, and the transient absorption spectra provide evidence for new transitions that were

not observed in the longer chain molecules. Molecular orbital calculations were carried out to explore these issues in more detail.

As was done in a previous Letter of peridinin [15], calculations were performed using a truncated chromophore, as shown in Figure 5. This molecule has C_s symmetry, and the higher symmetry and smaller size combine to make equation-of-motion coupled-cluster with singles and doubles (EOM-CCSD) calculations with polarizable continuum model (PCM) solvent effect theory tractable without changing any key properties of the π -system responsible for the spectroscopic properties [15]. The EOM-CCSD calculations presented in Figure 5 predict that the two lowest excited singlet states of C_{29} -peridinin are not only closer in energy than for peridinin, but strongly mixed, particularly in solvents with high refractive indices (see results for C_{29} -peridinin in tetrahydrofuran in Figure 5). The lowest two states in C_{29} -peridinin are more similar to the level ordering found in the protonated Schiff base retinal polyenes [29–32]. This prediction is in good agreement with the spectroscopic results.

A simulation of the transient absorption spectra of C_{29} -peridinin and peridinin in *n*-hexane and acetonitrile based on full single configuration interaction (CIS) methods and PCM solvent effect theory is presented in the left panel of Figure 6. The blue spectra were generated assuming the ground state geometry and that the lowest excited singlet state is the $2^1A_g^-$ state. The red spectra were generated based on the relaxed ICT (or $1^1B_u^+$ in *n*-hexane) geometry with all transitions occurring from that state. For spectral features involving C_{29} -peridinin, the lowest-lying relaxed singlet state in polar solvent is referred to as an ICT-like state, which should be taken to mean a weaker formation of the ICT state relative to that of peridinin (see below). Appropriately narrow bandwidths were used for the simulations so that the individual spectral features could be better discerned visually. The numbers at the top of each spectrum in Figure 6 indicate the index of the final singlet state. A comparison of the simulation (Figure 6) with the experimental

results (Figure 3) provides a useful perspective on the significant differences between C_{29} -peridinin and peridinin. While recognizing that a full CIS calculation will not be able to properly simulate the position or properties of the $2^1A_g^-$ state, the simulation does provide insights into the multiple bands that are present in the transient absorption spectra of C_{29} -peridinin.

The simulations of the transient absorption spectra shown in Figure 6 indicate that those of C_{29} -peridinin are quite different from those of peridinin. First, the intensities of the bands are less for all transitions in C_{29} -peridinin compared to analogous bands in peridinin. Second, there is one additional band predicted for both the set originating from the $1^1A_g^-$ state (blue spectra in Figure 6) and the relaxed $1^1B_u^+$ /ICT-like state (red spectra in Figure 6), which is consistent with the experimental observation of a long-lived band at 700 nm whose position does not depend on the solvent. The computations show that this additional band corresponds to the transition, S_1 (relaxed) $\rightarrow S_8$, and that it is more apparent in polar solvent due to separation from nearby bands that partially mask it. Moreover, the wavelength of the band is predicted to be insensitive to the solvent environment. Thus, the calculations are in good agreement with the general features of the observed spectra (Figure 3) and provide an explanation for the appearance of an additional band at 700 nm for C_{29} -peridinin. This band is also predicted to be present in peridinin in polar solvent (Figure 6), but it is significantly weaker and therefore not observed experimentally.

In view of these findings, it is important to reflect on the nature of the ICT state in C_{29} -peridinin and address the question of whether this molecule actually generates an ICT state or simply possesses a lowest-lying relaxed $1^1B_u^+$ state having the properties of an ICT state. The best perspective on this question is provided by the symmetry-adapted-cluster configuration-interaction (SAC-CI) calculations presented in the right panel of Figure 6 where the electrostatic properties of C_{29} -peridinin and peridinin in




Figure 6. Simulation of the transient absorption spectra (left panel) and comparison of the electrostatic properties (right panel) of C_{29} -peridinin and peridinin in various solvents. The results shown in the left panel were generated using full CIS methods to calculate the transition lengths, but with double excitation (CIS(D)) energy corrections for the origin states. The results shown in the right panel were generated using SAC-CI methods with level three selection of the doubly excited states.

methanol are shown for the relaxed excited singlet state. Examining the relaxed excited singlet state is critical to an analysis of the ICT state because both experiment and theory provide strong evidence that the ICT state is an evolved state that is created through relaxation in the singlet manifold [15]. In comparing the properties of the lowest-lying relaxed excited singlet state of C₂₉-peridinin with those of peridinin, significant differences observed when the molecules are dissolved in methanol (Figure 6) suggest that C₂₉-peridinin has an ICT-like state rather than a full ICT state for the following reasons: First, the dipole moment change for C₂₉-peridinin upon excitation into this state is significant ($\delta\mu = 8.4$ D), but it is a fraction of that calculated for peridinin ($\delta\mu = 21$ D). This value is considerable, but it is more similar to the ¹B_u⁺ states of polar retinyl polyenes and carotenoids than to the ICT state of peridinin [15,28–34]. Nevertheless, the dipole moment direction and charge shift contours shown in Figure 6 are indicative of a charge transfer from the allenic region into the lactone region; i.e., ICT-like behavior. Furthermore, one of the unique characteristics of the ICT state in peridinin is its significant doubly excited character (41.6%) [15], which is considerably larger than predicted for C₂₉-peridinin (19.8%). The smaller dipole moment and doubly excited character of the C₂₉-peridinin S₁ state is not due to a fundamental difference in the charge transfer process, rather it occurs as a consequence of the length of the polyene. A shorter polyene system cannot support as large a charge transfer configuration. Hence, it can be concluded that C₂₉-peridinin does indeed have an ICT state, but that this state is better viewed as an ICT-like or partial ICT state rather than a full ICT state as was found in the parent molecule, peridinin.

These findings reveal an apparent contradiction between the computations which indicate that C₂₉-peridinin does not support the formation of a full ICT state in polar solvents, and the experimental observations which reveal a large effect of solvent polarity on the lifetime of the lowest excited singlet state. The present investigation shows that it is an over-simplification to assume that the extent of ICT character is the sole factor controlling the dynamics of the lowest excited state of C₂₉-peridinin, peridinin, and other analogues when they are dissolved in polar solvents. This is because a short lifetime can also be associated with a level ordering reversal of the lowest ²1A_g⁻ and ¹1B_u⁺ states in polar solvents as is clearly the case for C₂₉-peridinin and peridinin in acetonitrile (Figure 5). Moreover, the excited state lifetimes of these molecules will scale inversely with the oscillator strengths of their relaxed singlet states [35]. Therefore, the impact of the ICT state on the lifetime of the lowest excited singlet state of peridinin and analogues is not due solely to its large dipole moment, but also to an increase in the oscillator strength of the ICT transition in polar solvents [15].

Acknowledgements

Work in the laboratory of HAF was supported by Grants from the National Science Foundation (MCB-1243565) and the University of Connecticut Research Foundation. Work in the laboratory of RRB was supported by Grants from the National Science Foundation (EMT-0829916), the National Institutes of Health (GM-34548) and the Harold S. Schwenk Sr. Distinguished Chair in Chemistry. Work in the laboratory of SK was supported by a Grant-in-Aid for Scientific Research on Innovative Areas 'Organic Synthesis based on Reaction Integration' and a Grant-in-Aid for Scientific

Research (C) 25410183 from the Ministry of Education, Culture, Sports, Science and Technology, Japan. Time-resolved spectroscopy was performed in the Photosynthetic Antenna Research Center (PARC), an Energy Frontier Research Center funded by the U.S. Department of Energy, Office of Science, Office of Basic Energy Sciences under Award Number DE-SC 0001035. HL was supported by an NSF Research Experience for Undergraduates (REU) grant to the Department of Chemistry at the University of Connecticut.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.cplett.2014.01.002>.

References

- [1] R.G. Hiller, in: H.A. Frank, A.J. Young, G. Britton, R.J. Cogdell (Eds.), *The Photochemistry of Carotenoids*, Kluwer Academic, Dordrecht, 1999, p. 81.
- [2] A.N. Macpherson, R.G. Hiller, in: B.R. Green, W.W. Parson (Eds.), *Light-Harvesting Antennas in Photosynthesis*, Kluwer Academic Publishers, 2002, p. 323.
- [3] M. Mimuro, S. Akimoto, in: A. Larkum, S. Douglas, J. Raven (Eds.), *Photosynthesis in Algae*, Kluwer Academic Publishers, 2003, p. 335.
- [4] J.A. Bautista et al., *J. Phys. Chem. B* 103 (1999) 8751.
- [5] H.A. Frank et al., *J. Phys. Chem. B* 104 (2000) 4569.
- [6] D. Zigmantas, R.G. Hiller, A. Yartsev, V. Sundström, T. Polivka, *J. Phys. Chem. B* 107 (2003) 5339.
- [7] D. Zigmantas, R.G. Hiller, F.P. Sharples, H.A. Frank, V. Sundström, T. Polivka, *Phys. Chem. Chem. Phys.* 6 (2004) 3009.
- [8] P. Chábera, M. Fuciman, P. Hřibek, T. Polivka, *Phys. Chem. Chem. Phys.* 11 (2009) 8795.
- [9] R. Pariser, *J. Chem. Phys.* 24 (1955) 250.
- [10] B. Hudson, B. Kohler, *Ann. Rev. Phys. Chem.* 25 (1974) 437.
- [11] P.R. Callis, T.W. Scott, A.C. Albrecht, *J. Chem. Phys.* 78 (1983) 16.
- [12] R.R. Birge, *Acc. Chem. Res.* 19 (1986) 138.
- [13] R.L. Christensen, E.A. Barney, R.D. Broene, M.G.I. Galinato, H.A. Frank, *Arch. Biochem. Biophys.* 430 (2004) 30.
- [14] R.L. Christensen, in: H.A. Frank, A.J. Young, G. Britton, R.J. Cogdell (Eds.), *The Photochemistry of Carotenoids*, Kluwer Academic Publishers, Dordrecht, 1999, p. 137.
- [15] N.L. Wagner, J.A. Greco, M.M. Enriquez, H.A. Frank, R.R. Birge, *Biophys. J.* 104 (2013) 1314.
- [16] T. Kajikawa et al., *Org. Lett.* 11 (2009) 5006.
- [17] D.M. Niedzwiedzki et al., *J. Phys. Chem. B* 113 (2009) 13604.
- [18] N. Chatterjee et al., *Arch. Biochem. Biophys.* 483 (2009) 146.
- [19] S. Kaligotla et al., *Photosynth. Res.* 103 (2010) 167.
- [20] D.M. Niedzwiedzki, T. Kajikawa, K. Aoki, S. Katsumura, H.A. Frank, *J. Phys. Chem. B* 117 (2013) 6874.
- [21] G. Britton, S. Liaaen-Jensen, H. Pfander, *Carotenoids Handbook*, Birkhäuser Verlag, Basel-Boston-Berlin, 2004.
- [22] D.M. Niedzwiedzki, M. Fuciman, H.A. Frank, R.E. Blankenship, *Biochim. Biophys. Acta, Bioenerg.* 1807 (2011) 518.
- [23] D. Zigmantas, T. Polivka, R.G. Hiller, A. Yartsev, V. Sundström, *J. Phys. Chem. A* 105 (2001) 10296.
- [24] F. Ehlers, D.A. Wild, T. Lenzer, K. Oum, *J. Phys. Chem. A* 111 (2007) 2257.
- [25] M. Kocpczynski, F. Ehlers, T. Lenzer, K. Oum, *J. Phys. Chem. A* 111 (2007) 5370.
- [26] I.H.M. van Stokkum, D.S. Larsen, R. van Grondelle, *Biochim. Biophys. Acta* 1657 (2004) 82.
- [27] D. Niedzwiedzki et al., *J. Phys. Chem. B* 111 (2007) 5984.
- [28] M.M. Enriquez, M. Fuciman, A.M. LaFountain, N.L. Wagner, R.R. Birge, H.A. Frank, *J. Phys. Chem. B* 114 (2010) 12416.
- [29] R. Mathies, L. Stryer, *Proc. Natl. Acad. Sci. U.S.A.* 73 (1976) 2169.
- [30] R.R. Birge, L.M. Hubbard, *J. Am. Chem. Soc.* 102 (1980) 2195.
- [31] R.R. Birge, C.F. Zhang, *J. Chem. Phys.* 92 (1990) 7178.
- [32] M. Ponder, R. Mathies, *J. Phys. Chem.* 87 (1983) 5090.
- [33] D. Zigmantas, R.G. Hiller, V. Sundström, T. Polivka, *Proc. Natl. Acad. Sci. U.S.A.* 99 (2002) 16760.
- [34] L. Premvardhan, E. Papagiannakis, R.G. Hiller, R. van Grondelle, *J. Phys. Chem. B* 109 (2005) 15589.
- [35] S.J. Strickler, R.A. Berg, *J. Chem. Phys.* 37 (1962) 814.